

Product Name: EuroChem SENTURA Fungicide  
APVMA Approval No: 87989 / 119975



Label Name:	EuroChem SENTURA Fungicide
-------------	----------------------------

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	--

Constituent Statements:	ACTIVE CONSTITUENT : 230 g/L Fludioxonil
-------------------------	--

Mode of Action:	GROUP 12 FUNGICIDE
-----------------	--------------------

Statement of Claims:	For the Control of certain post-harvest diseases in pome fruit as per the Directions for Use.
----------------------	---

Net Contents:	1L, 5L, 10L
---------------	-------------

Restrains:	
------------	--

Directions for Use:	This section contains file attachment.
---------------------	--

Other Limitations:	
--------------------	--

Withholding Periods:	NOT REQUIRED WHEN USED AS DIRECTED
----------------------	------------------------------------

Trade Advice:	
---------------	--

General Instructions:	This section contains file attachment.
-----------------------	--

Resistance Warning:	<p><b>GROUP 12 FUNGICIDE</b></p> <p>EuroChem SENTURA Fungicide is a member of the phenylpyrrole group of fungicides. For fungicide resistance management EuroChem SENTURA Fungicide is a Group 12 fungicide. Some naturally occurring individual fungi resistant to EuroChem SENTURA Fungicide and other Group 12 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by EuroChem SENTURA Fungicide and other Group 12 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, TGAC Australia Pty Ltd accepts no liability for any losses that may result from the failure of EuroChem SENTURA Fungicide to control resistant fungi.</p>
---------------------	---

Precautions:	Re-handling: DO NOT re-handle treated fruit until the product has been allowed to dry, unless wearing chemical resistant gloves.
--------------	--

Protections:	<p><b>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT</b></p> <p>Dangerous to fish and other aquatic organisms. DO NOT contaminate streams, rivers or waterways with the product or used containers.</p>
--------------	--

Storage and Disposal:	<p>Keep out of reach of children.</p> <p>Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.</p> <p>Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.</p> <p>If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p> <p><b>DIP DISPOSAL</b></p> <p>Unused or spent dip should be disposed of carefully to avoid contamination of streams, rivers or waterways. Dispose of dip in an authorised dip disposal facility.</p>
-----------------------	---

Safety Directions:	<p>May irritate the eyes. Harmful if inhaled. Repeated exposure may cause allergic disorders. Avoid contact with the eyes. DO NOT inhale spray mist.</p> <p>When opening the container and preparing the product for use, wear:</p> <ul style="list-style-type: none"> <li>• cotton overalls buttoned to the neck and wrist (or equivalent clothing)</li> </ul> <p>If applying by hand wear:</p> <ul style="list-style-type: none"> <li>• half face-piece respirator with organic vapour/gas cartridge or canister.</li> </ul> <p>Change cartridge if odour or taste of pesticide is noticed (a maximum of 8 hours of use is recommended).</p> <p>Wash hands after use. After each day's use, wash respirator and if rubber wash with detergent and warm water, and contaminated clothing.</p>
--------------------	--

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.
-------------------------	--

First Aid Warnings:	
---------------------	--

**DIRECTIONS FOR USE**

Crop	Disease	Rate		Critical Comments
		Dip and high volume	Low volume	
Pome Fruit	Blue Mould ( <i>Penicillium expansum</i> & <i>P. solitum</i> ), Grey Mould ( <i>Botrytis cinerea</i> )	130 to 260 mL in 100 L water	-	Refer to Application section below for application instructions.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

## **GENERAL INSTRUCTIONS**

**EuroChem SENTURA** is a protective fungicide used as a post-harvest treatment to control certain post-harvest diseases. **EuroChem SENTURA** may be applied as a post-harvest dip or drench in pome fruit.

**Note:** **EuroChem SENTURA** may be degraded by exposure to direct sunlight. Treated fruit should not be stored in direct sunlight.

### **Mixing**

Clean mixing and spraying equipment before use. Prepare no more treatment solution than is needed for the immediate operation. Shake the container before use to ensure uniform dispersion of contents before measuring quantity required. Half-fill the spray or mixing tank with clean water or wax/oil emulsion (or aqueous dilution of a wax/oil emulsion) and start agitation. While filling the remainder of the spray tank add the required amount of EuroChem SENTURA. Begin application of the solution after EuroChem SENTURA has completely and uniformly dispersed into the mix carrier. Maintain agitation throughout the treatment operation. DO NOT let the treatment mixture stand overnight in the tank. Flush the spray equipment thoroughly after each use.

### **Application**

#### ***Dip and High Volume Drench Application***

Dip fruit in prepared solution for 30 to 60 seconds and allow fruit to drain. Ensure fruit is in complete contact with the dipping solution.

### **Compatibility**

**EuroChem SENTURA Fungicide** can be mixed with Tecto®, imazalil (eg Fungiform®), calcium hypochlorite and DPA. A reduction in efficacy may occur when **EuroChem SENTURA** is applied with DPA at DPA rates above 500 ppm. If tank mixes are to be used observe all directions, precautions and limitations on all products to be used.

As formulations of other manufacturer's products are beyond the control of TGAC Australia Pty Ltd, and the quality of water may vary with location, all mixtures should be tested prior to mixing commercial quantities.

### **Cleaning Equipment**

Rinse application equipment with water after use. Spread rinsings over flat land away from desirable vegetation, waterways and drainage, and follow Dip Disposal instructions.